

Postural control performance of active and inactive older adults during manipulation of sensory information tests: a cross-sectional study

ESTELE C. W. M. LEMOS¹ | CLARISSA S. TEIXEIRA² | RAFAELA O. MACHADO³ | GABRIEL I. PRANKE⁴ | LUIZ F. C. LEMOS⁴ | CARLOS B. MOTA⁴

¹ Universidade Federal do Espírito Santo, Vitória, ES, Brazil

² Universidade Federal de Santa Catarina, Florianópolis, SC, Brazil

³ Pontifícia Universidade Católica do Paraná, Curitiba, PR, Brazil

⁴ Universidade Federal de Santa Maria, Santa Maria, RS, Brazil

Correspondence to: Estele Caroline Welter Meereis Lemos
Universidade Federal do Espírito Santo, ES, Brazil
ORCID 0000-0001-7827-1829
email: estelefisio@gmail.com
<https://doi.org/10.20338/bjmb.v18i1.410>

HIGHLIGHTS

- This study compared the postural balance of active and inactive elderly women in situations involving manipulation of sensory systems.
- This study corroborates the literature on the importance of physical exercise practice as a preventive strategy to reduce the occurrence of falls.
- Active elderly present reduced dependence of visual inputs and improved somatosensory afferences when compared to inactive elderly.
- Active elderly present higher COP average displacement speed than inactive elderly.

ABBREVIATIONS

ADL	Activities of daily living
ap	Anteroposterior
CNS	Central nervous system
COP	Center of pressure
COPap	COP displacement in the ap direction
COPml	COP displacement in the ml direction
COPvel	COP average displacement speed
Fap	Ground reaction force ap component
FLP	Foam-laser Dynamic Posturography method
Fml	Ground reaction force ml component
Fv	Ground reaction force vertical component
h	Distance from the surface to the geometric center of the platform
Map	Moment about the anteroposterior axis
ml	Mediolateral
Mml	Moment about the mediolateral axis
MMSE	Mini Mental State Examination
TOS	Sensory Organization Test

PUBLICATION DATA

Received 18 12 2023

Accepted 13 04 2024

Published 22 06 2024

BACKGROUND: Inactivity during the aging process negatively influences postural control.

AIM: The aim of the present study is to evaluate the postural balance of active and inactive elderly women in situations involving manipulation of sensory systems.

METHOD: The study included 20 elderly women (10 active and 10 inactive) with a mean age of 63.45 ± 2.37 years, who underwent postural balance assessment using a force platform in the six conditions of the Sensory Organization Test. For comparison between groups, the Student's t-test and Mann-Witney test were used. Statistically significant differences were observed between the groups for the three balance variables analyzed: Center of pressure displacement in the anteroposterior direction (COPap), in the mediolateral direction (COPml) and average speed (COPvel).

RESULTS: In situations in which the somatosensory system was not disturbed, as Sensory Organization Test (TOS) I, II and III, it was observed that active elderly women presented smaller COPap and COPml than inactive, however, in tests in which the somatosensory system was disturbed, due to the addition of a cushion between the platform and the individual's feet (TOS IV, V and VI), this difference was not observed. Furthermore, active elderly present higher COP displacement average speed than inactive elderly in almost all sensory tests (TOS I, III IV, V and VI).

CONCLUSION: It can be concluded that active elderly presented reduced dependence of visual inputs and improved somatosensory afferences when compared to inactive elderly and that active elderly presented higher COP average displacement speed than inactive elderly.

KEYWORDS: Postural balance | Elderly | Physical exercise

INTRODUCTION

Aging is characterized as a progressive and natural process with constant changes, biologically, functionally and structurally¹. It is a process that can alter the neuromuscular, somatosensory, vestibular and visual systems², which can influence postural stability,

generating deficits in the postural control system³.

In this context, it is known that balance is a complex function that requires the integration of the somatosensory, visual and vestibular systems, and the organization of this information by the central nervous system (CNS), which will generate neuromuscular responses^{4,5}. In this way, it can directly impact the activities of daily living (ADL) of the elderly^{3,6,7}. In the literature, there are studies that emphasize that these progressive changes delay the feedback of information, affecting postural control and consequently increasing the risk of falls^{8,9}.

Falls occur when the individual loses control of their center of gravity and there is insufficient recovery of balance¹⁰. In this way, they are characterized as any type of unintentional displacement to a level lower than the level before, with the inability to correct in a timely manner, even if a fall to the ground does not occur¹¹.

The causes of the increased risk of falls in the elderly can be multifactorial and subdivided into intrinsic, extrinsic and behavioral factors¹². Thus, it becomes a serious public health issue related to the age of individuals over 60 years old, as it has consequences such as increased risk of subsequent falls, hospital readmission, functional decline, mortality, fear of falling again and social isolation^{8,13}.

The practice of physical activity acts as an essential factor in maintaining functional fitness and postural balance, reducing the negative effects caused by the aging process, even delaying some restrictions regarding the performance of ADLs, prolonging the active life span of this population as well as reducing the rate of falls in the elderly population^{14,15,16}.

Studies have shown that physical inactivity harms the systems that act in postural control^{17,18} as well as that with the practice of physical activity there is an improvement in functional, coordinative and neuromotor capacity¹⁹. Furthermore, there are indicators that show that exercises improves postural and neuromotor stability, such as a decrease in the total speed of the trajectory and area of sway in the anteroposterior direction with eyes open and closed and bipedal support^{17,20}. However, it is still necessary to investigate which sensory systems physical exercise influences the most, so strategies and interventions can be proposed in order to focus on the system in question.

The amplitude and speed of body sway in the mediolateral and anteroposterior directions are frequently investigated COP variables^{9,21-23} and are used to predict the risk of falls in the elderly²⁴⁻²⁶ in addition to the sensory organization test, through assessment by Foam- laser Dynamic Posturography (FLP) method, which analyzes the relative contribution of the somatosensory, visual and vestibular systems to the individual's body stability, indicating the system responsible for instability, it is possible to infer that the use of the aforementioned tests can concomitantly bring interesting information about the means these systems contribute to postural control.

In this context, the aim of the present study was to evaluate the postural balance of active and inactive elderly women in situations involving manipulation sensorial to analyzes the relative contribution of sensory systems. Assessments of the visual, somatosensory and vestibular systems were carried out with eyes open and closed, with an unstable and stable surfaces and with and without a mobile cabin.

METHODS

This is a cross-sectional study, approved by the Ethics and Research Committee of the Federal University of Santa Maria (CAAE – 010.0.243.000-08), in accordance with the standards established in Resolution 466/12 of the National Health Council on research involving human beings.

Participants

The study included 20 elderly women who met the inclusion criteria: aged between 60 and 80 years, scoring above 26 points on the Mini Mental State Examination (MMSE), complying with the Informed Consent Form. The sample selection was for convenience, and for the group of active elderly women, elderly women who practice water aerobics at the Federal University of Santa Maria (UFSM) were invited, and those who performed physical exercises at a frequency equal to or greater than 150 minutes per week in the last three months²⁷ were included. For the group of inactive elderly women, elderly women were invited from a long-term care institution for elderly people living in Santa Maria-RS who had not performed physical exercise for 150 minutes per week in the last three months²⁸. Individuals who had known neurological disease, acute orthopedic trauma, severe visual impairment and/or used medications capable of influencing the tests were excluded from the study.

To calculate the sample size, G*Power software version 3.1.9.2 was used, in which an $\alpha = 0.05$ was adopted, a power of 90%. Of the variables to be investigated in the present study, the results found in the study by Meereis et al.¹⁶, referring to the amplitude of COP displacement in the mediolateral direction during the Sensory Organization Test (TOS 1) were used to support the sample calculation. Thus, the effect size was 1.42, therefore the minimum "n" of 10 individuals in each group was adopted.

Procedures

The research was carried out in the biomechanics laboratory of the Center for Physical Education and Sports (CEFD) at UFSM. First, an anamnesis and assessment of the level of physical activity²⁷ and MMSE were carried out to identify cognitive function. Soon after, anthropometric data such as body mass and height were collected using a scale and a Welmy stadiometer, respectively.

The group was composed of 10 active elderly women and 10 inactive elderly women, with similar anthropometric characteristics, illustrated in table 1.

Table 1. Anthropometric Characteristics.

	Active Elderly n= 10	Inactive Elderly n=10		
	Mean ± SD	Mean ± SD	T-Test	Mann-Whitney Test
Age (years)	61,35 ± 2,80	62,95 ± 2,87	0,183	
Height (m)	1,53 ± 0,12	1,54 ± 0,05		0,932
Mass (kg)	73,42 ± 7,03	75,34 ± 7,55	0,516	
BMI (kg/cm²)	29,56 ± 5,67	31,50 ± 3,63	0,585	

n= number of individuals in the group; BMI = body mass index; p-value = probability value in Student's t test, indicates a statistically significant difference if $p < 0.05$.

The assessment of postural control was carried out using a force platform (OR6-6, Advanced Mechanical Technology Incorporation, USA), during sensory manipulation, which was carried out through the sensory organization test (TOS), using posturography dynamics validated by Castagno²⁸. For the evaluation, each participant was positioned the force platform in a one m² cabin, two meters high.

Individuals were asked to remain standing barefoot on the platform during the six conditions following sensory organization test (TOS), with feet positioned hip-width apart for 30 seconds after the individual was stabilized. The support base was marked on each individual's first attempt on graph paper so that the positioning of the lower limbs could be repeated in all conditions. For unstable support surface conditions, an AIREX® Balance Pad (Airex AG, Inc.) flexible foam cushion was added on the platform with dimensions of 50 cm long, 50 cm wide and 10 cm thick and density, calculated by indications Patel²⁹, of 80 kg/m³.

Hereinafter the six conditions of the sensory organization test are presented:

- TOS I) Individual with eyes open and fixed horizontally, stable support surface (feet supported on the force platform), fixed visual cabin. This test evaluates the visual, somatosensory and vestibular systems.
- TOS II) Individual with eyes closed, stable support surface. Assessment of the somatosensory and vestibular systems.
- TOS III) Eyes open, stable foot support surface, oscillating visual cabin, 10 seconds manually tilted forward and 10 seconds to return to the starting position. It evaluates the somatosensory, vestibular and, above all, visual systems.
- TOS IV) Eyes open, fixed visual cabin, unstable support surface (with the addition of a foam cushion between the person's feet and the platform). It mainly evaluates the somatosensory system.
- TOS V) Eyes closed, fixed visual cabin, unstable support surface. This test evaluates the somatosensory and vestibular systems under overload conditions.
- TOS VI) Eyes open, unstable support surface with oscillating visual cabin. Assessment of the somatosensory, visual and vestibular systems.

Three attempts were made in each condition, with a 1-min interval between them, totaling 18 attempts. The order of tests was randomized for each participant. The displacement of the center of force overtime results in a trajectory which is related to the movement of the center of mass²¹ and, therefore, related with postural control. The COP position is calculated each time:

$$\text{COPap} = (\text{Mml} - h \cdot \text{Fap}) / \text{Fv}$$

$$\text{COPml} = (\text{Map} - h \cdot \text{Fml}) / \text{Fv}$$

Where COPap denotes center of pressure displacement in the anteroposterior direction; COPml denotes center of pressure displacement in the mediolateral direction; Map denotes moment about the anteroposterior axis; Mml denotes moment about the mediolateral axis; Fap: ground reaction force anteroposterior component; Fml denotes ground reaction force mediolateral component; Fv denotes ground reaction force vertical component, h denotes distance from the surface to the geometric center of the platform.

The variables analyzed were center of pressure displacement in the anteroposterior direction (COPap), in the mediolateral direction (COPml) and center of pressure average displacement speed (COPvel). The frequency used to acquire data from the force platform (forces and moments) was 100 Hz. The raw data from the platform were filtered by a 4th order Butterworth low-pass filter at a frequency of 10 Hz, thus eliminating possible interference from the environment.

Statistical Analyses

For statistical analysis, SPSS Software (Statistical Package for the Social Science) version 13.0 for Windows was used. To analyze the distribution of data, the Shapiro-Wilk test was performed. Then, the Student's t test was performed for data with normal distribution and the Mann-Whitney test for data with non-normal distribution for comparison between groups. For all tests, a significance level of 5% was used.

RESULTS

The results of the study are described below. Table 2 illustrates the behavior of COPap, COPml and COPvel during the execution of the six sensory organization tests.

Table 2. Balance test statistical data

		Active Elderly n= 10	Inactive Elderly n=10	T-Test	Mann-Whitney Test
		Mean ± SD	Mean ± SD		
TOS I	COPap (cm)	1,27 ± 0,69	2,74 ± 0,94		< 0,01*
	COPml (cm)	0,59 ± 0,30	1,70 ± 0,69	< 0,01*	
	COPvel (cm/s)	3,22 ± 1,70	1,26 ± 0,28		< 0,01*
TOS II	COPap (cm)	1,78 ± 0,74	2,59 ± 0,83	0,02*	
	COPml (cm)	1,00 ± 0,60	1,31 ± 0,51		< 0,01*
	COPvel (cm/s)	3,58 ± 1,44	1,40 ± 0,42		0,11
TOS III	COPap (cm)	2,38 ± 1,26	3,74 ± 1,21		0,03*
	COPml (cm)	0,94 ± 0,57	2,22 ± 1,04	< 0,01*	
	COPvel (cm/s)	3,43 ± 1,09	1,76 ± 0,64	< 0,01*	
TOS IV	COPap (cm)	2,14 ± 0,61	3,32 ± 1,77		0,03*
	COPml (cm)	1,24 ± 0,37	2,60 ± 2,49		0,10
	COPvel (cm/s)	3,64 ± 1,47	1,70 ± 0,72		< 0,01*
TOS V	COPap (cm)	3,54 ± 1,16	4,70 ± 1,69	0,05	
	COPml (cm)	1,97 ± 1,00	2,20 ± 0,86		0,35
	COPvel (cm/s)	4,19 ± 1,30	2,23 ± 0,67		< 0,01*
TOS VI	COPap (cm)	6,83 ± 5,75	6,41 ± 1,83		0,37
	COPml (cm)	3,37 ± 4,75	3,13 ± 1,11		0,26
	COPvel (cm/s)	5,38 ± 2,56	2,69 ± 0,96		< 0,01*

COPap = displacement of the center of pressure in the anteroposterior direction; COPml = displacement of the center of pressure in the medio-lateral direction, VM = Average velocity of COP displacement. * < 0.05.

DISCUSSION

Active elderly women demonstrated less postural sway, which infers better postural control and a lower risk of falls than inactive elderly women. By specifically analyzing the COP sway amplitude in different TOS conditions on a stable support surface, that is, TOS I, II and III, statistically significant differences were found for all postural balance variables analyzed. Indicating the contribution of the sensory systems responsible for postural balance without manipulating the somatosensory system.

In relation to TOS I, a situation in which information from the visual, vestibular and somatosensory systems were present, the COP displacement amplitude was significantly smaller in the group of active elderly women for both COPap and COPml, inferring that exercise promotes better postural control, corroborating other studies^{16,22}. Similarly, in TOS II and TOS III, when visual information is suppressed and respectively manipulated, the COPap and COPml displacement values from active elderly were lower than inactive. Elderly people generally depend more on visual inputs, and the results of this study indicated that active subjects presented better postural performance and managed sensory conflicting situations in a better way when compared to inactive subjects. Given this, it can be inferred that exercise can mitigate the risk of falls in situations where vision is compromised, such as, in situations of low vision or low light, circumstances which generate greater postural instability for the elderly³¹.

In situations of TOS IV, V and VI when an unstable support surface is added, hindering the action of the somatosensory system, the active elderly women obtained statistically significant lower values only in one of six variables (COPap on TOS IV). From this result, it can be observed that when somatosensory inputs were distorted, the sway amplitude of the two groups was similar. This finding corroborates other studies³² meaning that physical exercise improves somatosensory afferences, since it indicates that the active elderly group depend of undisturbed somatosensory inputs to present better balance control than inactive group.

Regarding COPvel, it can be observed that active elderly women presented statistically significant higher COP displacement speed in most tests (TOS I, TOS III, TOS IV, TOS V, TOS VI), differently from other studies that compared active and inactive elderly women^{22,36}. However, the present study suggests that active elderly women are able to make postural adjustments more quickly, ensuring the maintenance of postural control, demonstrating that the central nervous system was able to change the trajectory of sway quickly in order to ensure that balance was not threatened. This is important, since the ability to make postural adjustments necessary for adequate postural control is reduced with aging, especially in complex tasks^{9,37}.

Several authors have found a relationship between better postural control and physical exercise^{16,33,34}, showing that losses related to postural control are related not only to the aging process, but also to physical inactivity, thus, the practice of physical exercises is capable to improve the balance of inactive elderly women. In relation to this, Patti and collaborators³⁵, state that the practice of physical exercises favors the improvement of the conditions for receiving sensory information from the vestibular, visual and somatosensory systems and also activates the antigravity muscles, in addition to limiting the loss of muscle mass and strength, which is reinforced by the results of the present study.

This study as any cross-sectional analysis has certain limitation. The group of inactive elderly woman was institutionalized, so some characteristics arising from institutionalization can provide to these elderly more inactivity than living in community. However, the influence of institutionalization on elderly physical activity wasn't the aim of this study.

Despite the limitations, the present study reinforces that postural control is influenced by the practice of physical exercise and active elderly women have a smaller COP displacement compared to inactive elderly women^{9,38,39}. This reinforces the idea that the lower postural control and consequent higher risk of falls among inactive elderly women may not only be due to the aging process, but also to factors related to physical inactivity^{18,40}.

CONCLUSION

This study corroborates the literature on the importance of practicing physical exercise as a preventive strategy to reduce the occurrence of falls because (a) it demonstrates that active elderly present reduced dependence of visual inputs when compared to inactive elderly; (b) somatosensory afferences were improved and (c) higher COP average displacement speed can reflect the ability to make postural adjustments more quickly in order to maintain a postural control.

REFERENCES

1. Sgarbieri VC, Pacheco MTB. Healthy human aging: intrinsic and environmental factors. *Braz J Food Technol*. 2017;20:1-23. doi:10.1590/1981-6723.00717
2. Gschwind YJ, Kressig RW, Lacroix A, Muehlbauer T, Pfenninger B, Granacher U. A best practice fall prevention exercise program to improve balance, strength/power, and psychosocial health in older adults: study protocol for a randomized controlled trial. *BMC geriatrics*. 2013;13:1-13. doi:10.1186/1471-2318-13-105
3. Baltich J, Von Tscharnner V, Nigg BM. Degradation of postural control with aging. *Proc J Mech Eng H*. 2015;229(9):638-644. doi:10.1177/0954411915596013

4. Gerdhem P, Ringsberg KA, Akesson K. The relation between previous fractures and physical performance in elderly women. *Arch Phys Med Rehabil*. 2006;87:914-917. doi: 10.1016/j.apmr.2006.03.019
5. Duarte M, Freitas SM. Revisão sobre posturografia baseada em plataforma de força para avaliação do equilíbrio. *Braz J Phys Ther*. 2010;14(3):183-192. doi:10.1590/S1413-35552010000300003
6. Tieland M, Trouwborst I, Clark BC. Skeletal muscle performance and ageing. *J Cachexia Sarcopenia Muscle*. 2018;9:3–19. doi:10.1002/jcsm.12238
7. Gorgoulis V, Adams PD, Alimonti A, Bennett DC, Bischof O, Bishop C, et al. Cellular Senescence: Defining a Path Forward. *Cell*. 2019;179:813–827. doi:10.1016/j.cell.2019.10.005
8. Salari N, Darvishi N, Ahmadianah M, Shohaimi S, Mohammadi M. Global prevalence of falls in the older adults: A comprehensive systematic review and meta-analysis. *J Ort Sur. Res*. 2022;17:1-13. doi:10.1186/s13018-022-03222-1
9. Magalhães GV, Razuk M, Vieira LA, Rinaldi NM. Postural control performance of active and inactive older adults assessed through postural tasks with different levels of difficulty. *Motriz: Rev Educ Fis*. 2022;28:1-11. doi:10.1590/S1980-657420220015421
10. Papa EV, Hassan M, Bugnariu N. The Effects of Performance Fatigability on Postural Control and Rehabilitation in the Older Patient. *Curr Geriatr Rep*. 2016;5(3):172-178. doi:10.1007/s13670-016-0179-4
11. World Health Organization. *Ageing, & Life Course Unit. WHO global report on falls prevention in older age*. WHO. 2008.
12. World Health Organization: *Step Safely: Step safely: strategies for preventing and managing falls across the life-course*. Geneva: WHO. 2021.
13. Cameron ID, Dyer SM, Panagoda CE, Murray GR, Hill KD, Cumming RG, Kerse N. Interventions for preventing falls in older people in care facilities and hospitals. *Cochrane Database Syst Rev*. 2018;9:1-332. doi:10.1002/14651858.CD005465.pub4
14. Sherrington C, Michaleff ZA, Fairhall N, Paul SS, Tiedemann A, Whitney J, et al. Exercise to prevent falls in older adults: an updated systematic review and meta-analysis. *Bri J Sports Med*. 2017;51(24):1750-1758. doi:10.1136/bjsports-2016-096547
15. Muiños M, Ballesteros S. Does physical exercise improve perceptual skills and visuospatial attention in older adults? A review. *Eur Rev Aging Phys Act*. 2018;15(2):1-12. doi: 10.1186/s11556-018-0191-0
16. Meereis ECW, Favretto C, Souza JD, Gonçalves MP, Mota CB. Influence of hydrokinetic therapy in postural balance of institutionalized elderly people. *Motriz: Rev Ed Fis*. 2013; 19:269–277. doi:10.1590/s1980-65742013000200004
17. Avelar BP, Costa JNDA, Safons MP, Dutra MT, Bottaro M, Gobbi S, et al. Balance Exercises Circuit improves muscle strength, balance, and functional performance in older women. *AGE*. 2016;38(14):1-11. doi:10.1007/s11357-016-9872-7
18. Oliveira MR, Rubens AS, Dascal JB, Teixeira DC. Effect of different types of exercise on postural balance in elderly women: a randomized controlled trial. *Arch Gerontol Geriatr*. 2014;59(3):506-514. doi:10.1016/j.archger.2014.08.009
19. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee ML. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports and Exerc*. 2011;43(7):1334-1359. doi:10.1249/MSS.0b013e318213febf
20. Navega MT, Furlanetto MG, Lorenzo DM, Morcelli MH, Tozim BM. Effect of the Mat Pilates method on postural balance and thoracic hyperkyphosis among elderly women: a randomized controlled trial. *Rev Bras Geriatria e Ger*. 2016;19:465–472. doi:10.1590/1809-98232016019.150022
21. Lemos LFC, Ribeiro JS, Mota CB. Correlations between the center of mass and the center of pressure in active elderly people. *Rev Bras Cien Mov*. 2019;23:31-39. doi: 10.18511/0103-1716/rbcm.v23n1p31-39.
22. Santos PM, Pivetta HMF, Mota CB. Relationship between postural control and physical activity level in a group of elderly people. *Rev Bras Fis Ex*. 2015;14:50-55. doi:10.33233/rbfe.v14i1.132
23. Dorneles PP, Meereis ECW, Pranke GI, Mota CB. Relationship between plantar arch index and postural balance. *Rev Bras Cien Mov*. 2014;22:114-120. doi:10.18511/rbcm.v22i2.4583
24. Stel VS, Smit JH, Pluijm SM, Lips P. Balance and mobility performance as treatable risk factors for recurrent falling in older persons. *J Clin Ep*. 2003;56(7):659-668. doi:10.1016/s0895-4356(03)00082-9
25. Quijoux F, Nicolai A, Chairi I, Bargiotas I, Ricard D, Yelnik A, et al. A review of center of pressure (COP) variables to quantify standing balance in elderly people: Algorithms and open-access code. *Phys Rep*. 2021;9:1-32. doi:10.14814/phy2.15067
26. Ronda J, Galvañ B, Moneris E, Ballester F. Asociación entre síntomas clínicos y resultados de la posturografía computarizada dinámica. *Acta Otor Esp*. 2002;53(4):252-255. doi: 10.1016/S0001-6519(02)78308-8
27. Piercy KL, Troiano RP, Ballard RM, Carlson SA, Fulton JE, Galuska DA, et al. The Physical Activity Guidelines for Americans. *JAMA*. 2018;320(19):2020-2028. doi:10.1001/jama.2018.14854
28. Castagno LA. A new method for sensory organization tests: the foam- laser dynamic posturography. *Rev Bras Otor*. 1994;60(4):287-96. URL: oldfiles.bjorl.org/conteudo/acervo/acervo_english.asp?id=2600
29. Piasecki M, Ireland A, Jones DA, McPhee JS. Age-dependent motor unit remodelling in human limb muscles. *Biogerontology*. 2016;17(3):485-496. doi:10.1007/s10522-015-9627-3
30. Patel M, Fransson P A, Lush D, Gomez S. The effect of foam surface properties on postural stability assessment while standing. *Gait Posture*. 2008;28(4):649–656. doi:10.1016/j.gaitpost.2008.04.018
31. Chen EW, Amy S N F, Chan KM, Tsang WWN. Balance control in very old adults with and without visual impairment. *Eur J Appl Physiol*. 2012;12(5):1631-1636. doi:10.1007/s00421-011-2139-1

32. Buatois S, Gauchard GC, Aubry C, Benetos A, Perrin P. Current Physical Activity Improves Balance Control during Sensory Conflicting Conditions in Older Adults. *Sports Med.* 2007; 28(1):53-58. Doi: 10.1055/s-2006-924054
33. Low DC, Walsh GS, Arkesteijn M. Effectiveness of Exercise Interventions to Improve Postural Control in Older Adults: A Systematic Review and Meta-Analyses of Centre of Pressure Measurements. *Sports Med.* 2017;47:101-112. doi:10.1007/s40279-016-0559-0
34. Medeiros I, Pereira L, Salerno V, Gomes D. Effect of physical exercise on aging: differences in physical abilities between active and sedentary elderly people. *JIM.* 2022;3:49-61. doi:10.29073/jim.v3i1.591
35. Patti A, Zangla D, Sahin FN, Cataldi S, Lavanco G, Palma A, Fischietti F. Physical exercise and prevention of falls. Effects of a Pilates training method compared with a general physical activity program: A randomized controlled trial. *Med Bal.* 2021;100(13):1-7. doi: 10.1097/MD.00000000000025289
36. Victor LGV, Oliveira MRD, Teixeira DDC, Paes MA, Fujisawa DS, Bispo NDND, et al. Postural control during one-leg stance in active and sedentary older people. *Motriz: Rev Ed Fis.* 2014;20:339–345. doi:10.1590/s1980-65742014000300014
37. Gotardi GC, Rodrigues ST, Godoi D, Barbieri F, Navarro M, Schor P, et al. The effects of horizontal and vertical saccadic eye movements on postural control of young adults and elderly in different bases of support. *Rev Bras Educ Fis Esporte.* 2018;32(4):559-68. doi: 10.11606/1807-5509201800040559
38. da Costa JL, Porolnik S, Arruda GT, Petter GN, Piveta HMF. Fall risk in physically active elderly: a cross-sectional study. *Braz J Dev.* 2021;7(1):6028–6037. doi:10.34117/bjdv7n1-410
39. Ceccon FG, CARPES FP. Implications of regular exercise on postural control in the elderly. *Interdisciplinary Studies on Aging.* 2015;20:139-158. doi: 10.22456/2316-2171.45275
40. Sarto F, Pizzichemi M, Chiossi F, Bisiacchi PS, Franchi MV, Narici MV, et al. Physical active lifestyle promotes static and dynamic balance performance in young and older adults. *Front Phys.* 2022;13:1-7. doi:10.3389/fphys.2022.986881

Citation: Meereis-Lemos ECW, Teixeira CS, Machado RO, Pranke GI, Lemos LFC, Mota CB. (2024). Postural control performance of active and inactive older adults during manipulation of sensory information tests: a cross-sectional study. *Brazilian Journal of Motor Behavior*, 18(1):e410.

Editor-in-chief: Dr Fabio Augusto Barbieri - São Paulo State University (UNESP), Bauru, SP, Brazil.

Associate editors: Dr José Angelo Barela - São Paulo State University (UNESP), Rio Claro, SP, Brazil; Dr Natalia Madalena Rinaldi - Federal University of Espírito Santo (UFES), Vitória, ES, Brazil; Dr Renato de Moraes - University of São Paulo (USP), Ribeirão Preto, SP, Brazil.

Guest editors: Dr Paula Favaro Polastri Zago - São Paulo State University (UNESP), Bauru, SP, Brazil; Dr Daniela de Godoi Jacomassi - Federal University of São Carlos (UFSCAR), São Carlos, SP, Brazil.

Copyright: © 2024 Lemos, Teixeira, Machado, Pranke, Lemos and Mota and BJMB. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives 4.0 International License which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Nothing to declare.

Competing interests: The authors have declared that no competing interests exist.

DOI: <https://doi.org/10.20338/bjmb.v18i1.410>